

# The nutritional value of *Moringa oleifera* Lam. leaves: what can we learn from figures?

Mélanie BROIN – Scientific Assistant – Moringanews Network ([melanie.broin@wanadoo.fr](mailto:melanie.broin@wanadoo.fr))

## Introduction

There are quite a lot of data available in the literature, on the Internet and from other sources about the nutritional value of *Moringa oleifera* Lam. (M.o.) leaves. However, it can sometimes be quite hard to find one's way through, since there can be considerable variation amongst data. In this context, we thought it might be useful for fieldworkers and for efficient communication about Moringa, to have reliable average figures on the nutritional value of M.o. leaves. The objective of this work was to gather available information about M.o. leaves nutritional value, to identify potential sources of errors in data, to discard irrelevant data, and to find a consensus on average nutritional values.

## Results

### 1. Source data

In total, we collected 23 references dealing with M.o. nutritional values, 11 of them were from peer reviewed journals, 5 from books or reports, 5 from unpublished analyses, and 2 from an Internet source (FAO).

### 2. Variability of source data

The ratio of the maximal to the minimal value in initial data varied from 1.9 to 514 (Table 1), illustrating the very high variability of source data. Furthermore, the high variability of source data is illustrated by Figure 1, which shows the standard deviation for each nutrient.

### 3. Selection of relevant data

Table 1 shows the ratios of maximal to minimal values obtained after having discarded outliers (the number of figures kept is indicated in parentheses). Figure 1 illustrates the remaining variability of selected data, by showing the standard deviation for each nutrient.

Table 1: Ratio of maximum to minimum value for the whole set of data (left column) and for the selected set of data (right column). The figures in parentheses indicate the number of data considered.

\*Vitamin C: values for fresh leaves only.

Nutrient	Ratio Maximum Value/Minimum Value	
	Source data	Selected data
Minerals	1.9 (9)	1.9 (9)
Manganese	1.9 (7)	1.9 (7)
Carbohydrates	4 (8)	1.6 (6)
Vitamin C*	4 (8)	0.8 (5)
Fibers	6 (8)	1.7 (4)
Vitamin E	8 (3)	N/A
Calcium	19 (16)	1.7 (13)
Proteins	24 (15)	1.8 (11)
Fat	25 (11)	1.9 (5)
Copper	29 (10)	2.2 (8)
Iron	30 (14)	2.1 (10)
Vitamin A	42 (10)	2.9 (7)
Phosphorus	64 (11)	1.5 (5)
Magnesium	76 (9)	1.4 (6)
Vitamin B2	111 (3)	N/A
Zinc	121 (10)	2.1 (7)
Vitamin B1	340 (4)	N/A
Potassium	514 (8)	2.2 (6)




Figure 1: Representation of the variability of data. Means for all nutrients have been set to 100 arbitrary units. Thin lines represent the standard deviation for the whole set of data, while boxes represent the standard deviation for the set of selected data. \*Vitamin C: values for fresh leaves only.

### 4. Possible sources of variation amongst data

Table 2 illustrates the possible sources of variation amongst data. While there is generally an acceptable range of variation due to differences in the genetic background, the environment, the cultivation methods and the sample analytical method, attention must be drawn on variations due to the methods of preparation and conservation of samples that can seriously affect the vitamin content. As far as errors of Human origin are concerned, they should be as far as possible identified and the values not taken into account.

Table 2: Possible sources of variations amongst data	
<b>Natural variation amongst source samples</b>	
Genetic background (ecotype, cultivar)	Ex: calcium content varies from 1400 to 2300 mg/100g DM in the selected data
Environment (soil, climate, pathogens)	
Cultivation methods (inputs, frequency of harvests)	
<b>Variation due to sample preparation and analysis</b>	
Time between collection and analysis	Ex: vitamin C content is 40 times lower in dried leaves (around 20 mg/100g DM) compared to fresh leaves (around 800 mg/100g DM)
Mode of conservation between collection and analysis (drying, refrigeration, freezing, etc.)	
Analytical method	
<b>Errors of Human origin</b>	
Error of manipulation during analysis	Ex: the values for zinc vary between 0.5 and 3.6 mg/100g DM, except for one value which is 26 mg/100g DM (ref.2), while the average value is 2.5mg/100g DM.
Error in the calculation of results	
Error in the edition of results	Ex. in ref.21, data reported for 100 mg dried leaves are: proteins=72mg; calcium=690 mg; iron=3.5mg; potassium=3.8mg; zinc=0.2mg; copper=0.15mg; vitamin C=1.1mg. These values are completely inconsistent with average values.
Error in the botanical identification of the sample	

## Methodology


We did an exhaustive search for data on the nutritional value of M.o. leaves through literature databases, the Internet, and the Moringanews Network. All the data collected were compiled in a Microsoft Excel spreadsheet for further analyses. When there were several analyses in one reference, they were compiled in order to give the same weight to each reference. All the data collected, whether from fresh or dry leaves, were transformed to obtain a value for 100g of dry matter, using the water content given in the analysis if available, or using the average water content for fresh or dry leaves, accordingly, if not. Data for vitamin A were all converted to International Units (IU) using a conversion factor beta-carotene to retinol of 1/6 and 1IU=0.3µg retinol, except in Figure 3 where a conversion factor of 1/18 has been used to avoid overestimation. For vitamin C, only values for fresh leaves have been considered, since there is a loss of vitamin C when the leaves are dried, except in Figure 3 where we used the value for dry leaves, since this Figure is dealing with leaf powder. We then calculated, for each nutrient, the mean, the standard deviation and the ratio of the maximal to the minimal value. In order to eliminate irrelevant values, we discarded outliers for each nutrient. We then recalculated the mean and the standard deviation for the data kept. So obtained mean values were then compared to other foods and to the recommended dietary allowances. We also compared the discarded data to the mean values to try to understand the potential causes of such high variations or errors.

### 5. Comparison of mean nutritional values with other foods and RDA

Table 3 shows, for each nutrient for which sufficient data were available, the mean values obtained and the standard deviation.

Figure 2 shows the nutritional interest of M.o. leaves in country where the diet is deficient in proteins, vitamins and minerals. Fresh M.o. leaves contain at least twice more proteins than milk, and half the proteins of eggs. They are richer in iron than lentils and beef meat, richer in calcium than milk, at least as rich as carrots in vitamin A, and richer in vitamin C than oranges.

But Figure 2 also highlights that Moringa is not a 'miracle food', as similar properties can be found in other leafy vegetables like cassava or amaranth leaves.



Nutrient	Mean value for 100g Dry Matter
Proteins (g)	29 ± 6
Minerals (g)	11 ± 2.2
Fat (g)	8 ± 2.5
Carbohydrates (g)	38 ± 7
Fibers (g)	10 ± 3
Calcium (mg)	1924 ± 288
Copper (mg)	1.0 ± 0.2
Iron (mg)	28 ± 6
Potassium (mg)	1384 ± 420
Magnesium (mg)	422 ± 52
Phosphorus (mg)	267 ± 49
Manganese (mg)	8.4 ± 2.4
Zinc (mg)	2.5 ± 0.6
Vitamin A (IU)	15620 ± 6475
Vitamin C* (mg)	773 ± 91

Table 3: Mean nutritional values of *Moringa oleifera* leaves. Data are given for 100g dry matter. \*Data originate from the analysis of both fresh and dry leaf samples, except for vitamin C for which only analyses of fresh samples have been taken into account, due to the loss of vitamin C during the drying process.

Figure 2: Comparison of the average nutritional values of *Moringa oleifera* fresh leaves with other foods. For M.o., the ellipse represents the range of acceptable values (mean ± SD). For other foods, the circle indicates the mean value from Food Composition Table for Use in Africa, FAO, 1968.

Figure 3 highlights the interest of *Moringa oleifera* leaf powder to prevent malnutrition in developing countries, that mostly appear in children during the weaning period, between 1 and 3 years old. Indeed, for children under three, 30g of leaf powder can cover one third of the daily allowance for proteins, 75% of the calcium needs, more than half of the iron necessary, the totality of the recommended dietary allowance for vitamin A, and almost one third of the needs in vitamin C. The leaf powder is also an interesting dietary supplement for pregnant and lactating women.




Figure 3: Percentage of coverage of recommended dietary allowances with 30g *Moringa oleifera* leaf powder, for children 1-3 years old, pregnant and lactating women. RDA are from *La nutrition dans les pays en développement*, Michael C. Latham, FAO, 2001, for a diet poor in proteins from animal sources and poor in vitamin C. \*For vitamin A, to avoid overestimation, a conversion rate beta-carotene to retinol of 1/18 has been used instead of 1/6 generally used, since 1/18 is closer to the real conversion rate for leafy vegetables. \*\*Value for vitamin C originates from only two sources from the analysis of leaf powder (data from fresh leaves have not been taken into account since they would be overestimated for leaf powder).

## References

- Booth and Wickens. 1988. Non-timber uses of selected wild trees and shrubs in Africa. FAO Conservation Guide, Rome, pp. 92-101. (Fresh leaves)
- Barnes et al. 1998. Mineral composition of non-conventional leafy vegetables. Plant Foods for Human Nutrition 53, pp. 29-36. (Fresh leaves)
- Freiberger et al. 1998. Nutrient content of the edible leaves of seven wild plants from Niger. Plant Foods for Human Nutrition 53, pp. 57-69. (Fresh leaves)
- Szefermala et al. 1998. Effect of cooking on the nutritive value of common food plants of Tanzania. Part 1-Vitamin C in some of the wild green leafy vegetables. Food Chemistry 10, pp. 205-210. (Fresh leaves)
- Chang. 2000. Alpha-tocopherol content in 62 edible tropical plants. Journal of Agriculture and Food Chemistry 49, pp. 3101-3105. (Fresh leaves)
- Hamply et al. 2002. Food diversity and drought survival: The Heuza example. International Journal of Food Science and Nutrition 44, pp. 1-16. (Fresh leaves)
- Coole et al. 2003. The distribution, uses and potential for development of *Moringa oleifera* Lam. in Malawi. Forestry Research Record 67, Forestry Research Institute of Malawi, Zomba. (Fresh leaves)
- D'Souza and Kulkarni. 2004. Comparative studies on nutritive values of tender foliage of seedlings and mature plants of *Moringa oleifera* Lam. Journal of Economic and Taxonomic Botany. (Fresh leaves)
- Ruby et al. 2000. Chemical composition of fruits and leaves eaten by short-nosed fruit bat, *Cynopterus sphinx*. Journal of Chemical Ecology 26, pp. 2825-2841. (Fresh leaves)
- Lokite et al. 1997. Energy and macromineral composition of dietary and medicinal wild plants consumed during drought. Study of rural Punjab, Northeastern Nigeria. International Journal of Food Science and Nutrition 51, pp. 195-206. (Dry leaves)
- Lokite et al. 1997. Tenure and conventional Ache foodies series: tradicões e alimentos do Níger. Cahiers Agricultures 6, pp. 933-940. (Dry leaves)
- Sera et al. 1998. Analysis of nutritional components of eight leafy foods of the Niger. Cahiers Agricultures 6, pp. 933-940. (Dry leaves)
- FAO. 2003. Analysis of 800. Montpellier, France. (Dry leaves)
- Binga trees. 2005. Analysis. (Dry leaves)
- FAO. 2004. Analysis. University of Zimbabwe. (Dry leaves)
- FAO. 1968. Food composition table for use in Africa. (Dry leaves)
- Tree Africa. 2006. Analysis. (Dry leaves)